

ROM2F-97/38
hep-th/9711077

Some Properties of Tensor Multiplets in Six-Dimensional Supergravity

Fabio Riccioni¹ and Augusto Sagnotti

*Dipartimento di Fisica
Università di Roma “Tor Vergata”
I.N.F.N. - Sezione di Roma “Tor Vergata”
Via della Ricerca Scientifica, 1
00133 Roma ITALY*

Abstract

We review some results on the complete coupling between tensor and vector multiplets in six-dimensional $(1,0)$ supergravity.

Contribution to the Proceedings of the String Duality II Conference, Trieste, ITALY, april 1997

¹I.N.F.N. Fellow

Some Properties of Tensor Multiplets in Six-Dimensional Supergravity

F. Riccioni[†] and A. Sagnotti

Dipartimento di Fisica, Università di Roma “Tor Vergata”

I.N.F.N., Sezione di Roma “Tor Vergata”

Via della Ricerca Scientifica, 1 00133 Roma ITALY

We review some results on the complete coupling between tensor and vector multiplets in six-dimensional $(1, 0)$ supergravity.

1. Minimal Six-Dimensional Supergravity with Vector and Tensor Multiplets

In the low-energy supergravity, the Green-Schwarz mechanism [1] results from couplings between antisymmetric tensors and vector fields whose anomalous behavior, for particular choices of the gauge group, cancels the reducible portion of the anomaly generated by fermion loops. In ten dimensions, this restricts the possible non-abelian groups to two choices, $SO(32)$ or $E_8 \times E_8$, while the residual, factorized anomaly is disposed of by the single antisymmetric tensor of the supergravity multiplet. On the other hand, in six dimensions the irreducible part of the gravitational anomaly cancels only provided [2]

$$n_H - n_V + 29n_T = 273 \quad , \quad (1)$$

where n_V , n_T and n_H are the numbers of vector, tensor and hypermultiplets, a looser constraint compatible with a large number of non-abelian gauge groups. Moreover, in this case one can naturally build type-I models with variable numbers of tensor multiplets [3] as parameter-space orbifolds (orientifolds) [4] of $K3$ reductions of the type-IIB string, where several antisymmetric tensors take part in the anomaly cancellation [5]. The two cases show another marked difference. While the ten-dimensional Green-Schwarz coupling, $B \wedge (F^4 - R^4)$, is a higher derivative term not visible in the low-energy lagrangian, the gauge portion of the corresponding six-dimensional coupling, $B \wedge (F^2 - R^2)$, has sizeable effects on the low-energy dynamics.

This talk is devoted to $(1, 0)$ supergravity in six dimensions coupled to n tensor multiplets and to vector multiplets. It reflects only in part the lecture presented by one of us (A.S.) at the “String Duality II” Conference, since it expands the portion of the original discussion devoted to tensor multiplet couplings in six-dimensional $(1, 0)$ supergravity. For completeness, we are also including some new results, that are fully presented in [6].

The starting point is the coupling of $(1, 0)$ supergravity to n tensor multiplets, originally studied by Romans [7] to lowest order in the fermi fields. Nishino and Sezgin [8] considered the coupling to a single tensor multiplet and to vector and hypermultiplets to all orders in the fermi fields, and have recently included some higher-order fermionic couplings in the theory with tensor, vector and hypermultiplets [9]. Here we complete the field equations and relate them to Wess-Zumino consistency conditions, in the spirit of [10], for the case of an arbitrary number of tensor and vector multiplets. The n scalars in the tensor multiplets parameterize the coset space $SO(1, n)/SO(n)$, and are described by the $SO(1, n)$ matrix [7]

$$V = \begin{pmatrix} v_r \\ x_r^m \end{pmatrix} \quad . \quad (2)$$

All spinors are symplectic Majorana-Weyl. In particular, the tensorini χ^m are right-handed, while the gravitino Ψ_μ and the gaugini λ are left-handed. The tensor fields $B_{\mu\nu}^r$ are valued in the fundamental representation of $SO(1, n)$, and their field strengths include Chern-Simons 3-forms of

[†]I.N.F.N. Fellow

the vector fields according to

$$H^r = dB^r - c^{rz}\omega_z \quad , \quad (3)$$

where the c^{rz} are constants that determine the gauge part of the residual anomaly polynomial

$$\mathcal{A} = \sum_{rs} \eta_{rs} c^{rx} c^{sy} tr_x(F \wedge F) tr_y(F \wedge F) \quad (4)$$

and z runs over the various factors of the gauge group [5]. Gauge invariance of H^r then requires that

$$\delta B^r = c^{rz} tr_z(\Lambda dA) \quad . \quad (5)$$

To lowest order in the fermi fields, the fermionic equations are [5]

$$\begin{aligned} & \gamma^{\mu\nu\rho} D_\nu \Psi_\rho + v_r H^{r\mu\nu\rho} \gamma_\nu \Psi_\rho \\ & - \frac{i}{2} x_r^m H^{r\mu\nu\rho} \gamma_{\nu\rho} \chi^m \\ & + \frac{i}{2} x_r^m \partial_\nu v^r \gamma^\nu \gamma^\mu \chi^m \\ & - \frac{1}{\sqrt{2}} v_r c^{rz} tr_z(F_{\sigma\tau} \gamma^{\sigma\tau} \gamma^\mu \lambda) = 0 \quad , \end{aligned} \quad (6)$$

$$\begin{aligned} & \gamma^\mu D_\mu \chi^m - \frac{1}{12} v_r H^{r\mu\nu\rho} \gamma_{\mu\nu\rho} \chi^m \\ & - \frac{i}{2} x_r^m H^{r\mu\nu\rho} \gamma_{\mu\nu} \Psi_\rho \\ & - \frac{i}{2} x_r^m \partial_\nu v^r \gamma^\mu \gamma^\nu \Psi_\mu \\ & - \frac{i}{\sqrt{2}} x_r^m c^{rz} tr_z(F_{\mu\nu} \gamma^{\mu\nu} \lambda) = 0 \end{aligned} \quad (7)$$

and

$$\begin{aligned} & v_r c^{rz} \gamma^\mu D_\mu \lambda + \frac{1}{2} (\partial_\mu v_r) c^{rz} \gamma^\mu \lambda \\ & + \frac{1}{2\sqrt{2}} v_r c^{rz} F_{\alpha\beta} \gamma^\mu \gamma^{\alpha\beta} \Psi_\mu \\ & + \frac{i}{2\sqrt{2}} x_r^m c^{rz} F_{\mu\nu} \gamma^{\mu\nu} \chi^m \\ & - \frac{1}{12} c^{rz} H_{r\mu\nu\rho} \gamma^{\mu\nu\rho} \lambda = 0 \quad , \end{aligned} \quad (8)$$

while the bosonic equations are

$$\begin{aligned} & R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R + \partial_\mu v^r \partial_\nu v_r \\ & - \frac{1}{2} g_{\mu\nu} \partial_\alpha v^r \partial^\alpha v_r - G_{rs} H_{\mu\alpha\beta}^r H^s{}_\nu{}^{\alpha\beta} \\ & + 4v_r c^{rz} tr_z(F_{\alpha\mu} F^\alpha{}_\nu) \\ & - \frac{1}{4} g_{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} = 0 \quad , \end{aligned} \quad (9)$$

$$\begin{aligned} & x_r^m \nabla_\mu (\partial^\mu v^r) + \frac{2}{3} x_r^m v_s H_{\alpha\beta\gamma}^r H^{s\alpha\beta\gamma} \\ & - x_r^m c^{rz} tr_z(F_{\alpha\beta} F^{\alpha\beta}) = 0 \end{aligned} \quad (10)$$

and

$$\nabla_\mu (v_r c^{rz} F^{\mu\nu}) - c^{rz} G_{rs} H^{s\nu\rho\sigma} F_{\rho\sigma} = 0 \quad . \quad (11)$$

Moreover, the tensor fields satisfy (anti)self-duality conditions, conveniently summarized as [10]

$$G_{rs} H^{s\mu\nu\rho} = \frac{1}{6e} \epsilon^{\mu\nu\rho\alpha\beta\gamma} H_{r\alpha\beta\gamma} \quad , \quad (12)$$

where $G_{rs} = v_r v_s + x_r^m x_s^m$. Under the supersymmetry transformations

$$\begin{aligned} & \delta e_\mu{}^a = -i(\bar{\epsilon} \gamma^a \Psi_\mu) \quad , \\ & \delta B_{\mu\nu}^r = i v^r \bar{\Psi}_{[\mu} \gamma_{\nu]} \epsilon + \frac{1}{2} x^{mr} \bar{\chi}^m \gamma_{\mu\nu} \epsilon \\ & - 2c^{rz} tr_z(A_{[\mu} \delta A_{\nu]}) \quad , \\ & \delta v_r = x_r^m \bar{\chi}^m \epsilon \quad , \\ & \delta A_\mu = -\frac{i}{\sqrt{2}} (\bar{\epsilon} \gamma_\mu \lambda) \quad , \\ & \delta \Psi_\mu = D_\mu \epsilon + \frac{1}{4} v_r H_{\mu\nu\rho}^r \gamma^{\nu\rho} \epsilon \quad , \\ & \delta \chi^m = \frac{i}{2} x_r^m \partial_\alpha v^r \gamma^\alpha \epsilon + \frac{i}{12} x_r^m H_{\alpha\beta\gamma}^r \gamma^{\alpha\beta\gamma} \epsilon \quad , \\ & \delta \lambda = -\frac{1}{2\sqrt{2}} F_{\mu\nu} \gamma^{\mu\nu} \epsilon \quad , \end{aligned} \quad (13)$$

the fermionic field equations turn into the bosonic ones, and this proves supersymmetry to lowest order.

This theory has a gauge anomaly, as one can see from the divergence of the (non-integrable) vector equation [10]

$$\begin{aligned} & \nabla_\mu J^\mu = \\ & - \frac{1}{2e} \epsilon^{\mu\nu\alpha\beta\gamma\delta} c^{rz} c_r^{z'} F_{\mu\nu} tr_{z'}(F_{\alpha\beta} F_{\gamma\delta}) \quad , \end{aligned} \quad (14)$$

that reproduces the residual covariant anomaly. An integrable equation,

$$\begin{aligned} & \nabla_\mu (v_r c^{rz} F^{\mu\nu}) - c^{rz} G_{rs} H^{s\nu\rho\sigma} F_{\rho\sigma} \\ & - \frac{1}{8e} \epsilon^{\nu\rho\alpha\beta\gamma\delta} c_r^{z'} A_\rho c^{rz'} tr_{z'}(F_{\alpha\beta} F_{\gamma\delta}) \\ & - \frac{1}{12e} \epsilon^{\nu\rho\alpha\beta\gamma\delta} c_r^z F_{\rho\alpha} c^{rz'} \omega_{\beta\gamma\delta}^{z'} = 0 \quad , \end{aligned} \quad (15)$$

defines a gauge current that generates the residual consistent anomaly. However, the resulting theory is no longer supersymmetric, but has a corresponding supersymmetry anomaly [10]

$$\mathcal{A}_\epsilon = -c^{rz}c_r^{z'}[tr_z(\delta_\epsilon AA)tr_{z'}(F^2) - 2tr_z(\delta_\epsilon AF)\omega_3^{z'}] \quad , \quad (16)$$

determined by the Wess-Zumino consistency conditions [11]. Moreover, one can prove that the divergence of the gravitino equation yields the supersymmetry anomaly.

As usual, the commutators of supersymmetry transformations close only on-shell on fermi fields. In order to complete the construction, *all* first order field equations and the supersymmetry transformations of the fermi fields must be made supercovariant. It should be appreciated that the former include the (anti)self-duality conditions (12) on tensor fields [12]. The complete theory can then be constructed imposing closure of the algebra on the fermi fields. This requires a long and tedious calculation, with the end result that the algebra closes on-shell in the expected fashion on all fermi fields, with the notable exception of the gaugini. Indeed, in this case the algebra generates all the local symmetries and the field equation, together with an additional term

$$\begin{aligned} [\delta_1, \delta_2]_{extra}\lambda = & \\ & \frac{c \cdot c'}{v \cdot c} tr_{z'}[-\frac{1}{4}(\bar{\epsilon}_1 \gamma_\alpha \lambda')(\bar{\epsilon}_2 \gamma_\beta \lambda')\gamma^{\alpha\beta} \lambda \\ & + \frac{1}{4}(\bar{\lambda} \gamma_\alpha \lambda')(\bar{\epsilon}_1 \gamma^\alpha \lambda')\epsilon_2 - (1 \leftrightarrow 2) \\ & + \frac{1}{16}(\bar{\epsilon}_1 \gamma^\alpha \epsilon_2)(\bar{\lambda}' \gamma_{\alpha\beta\gamma} \lambda')\gamma^{\beta\gamma} \lambda] \quad . \end{aligned} \quad (17)$$

We will return in the next Section to the crucial role of this charge.

From the complete fermionic equations, one can deduce a lagrangian that generates them, and finally derive from it the complete bosonic equations. The lagrangian is

$$\begin{aligned} e^{-1}\mathcal{L} = & -\frac{1}{4}R - \frac{1}{2}v_r c^{rz} tr_z(F_{\alpha\beta} F^{\alpha\beta}) \\ & + \frac{1}{12}G_{rs} H^{r\mu\nu\rho} H_{\mu\nu\rho}^s - \frac{1}{4}\partial_\mu v^r \partial^\mu v_r \\ & - \frac{1}{8e}\epsilon^{\mu\nu\alpha\beta\gamma\delta} c_r^z B_{\mu\nu}^r tr_z(F_{\alpha\beta} F_{\gamma\delta}) \end{aligned}$$

$$\begin{aligned} & -\frac{i}{2}\bar{\Psi}_\mu \gamma^{\mu\nu\rho} D_\nu \Psi_\rho - \frac{i}{2}v_r H^{r\mu\nu\rho}(\bar{\Psi}_\mu \gamma_\nu \Psi_\rho) \\ & + \frac{i}{2}\bar{\chi}^m \gamma^\mu D_\mu \chi^m - \frac{i}{24}v_r H_{\mu\nu\rho}^r(\bar{\chi}^m \gamma^{\mu\nu\rho} \chi^m) \\ & + \frac{1}{2}x_r^m \partial_\nu v^r (\bar{\Psi}_\mu \gamma^\nu \gamma^\mu \chi^m) \\ & - \frac{1}{2}x_r^m H^{r\mu\nu\rho}(\bar{\Psi}_\mu \gamma_{\nu\rho} \chi^m) \\ & + \frac{i}{\sqrt{2}}\bar{\Psi}_\mu \gamma^{\sigma\delta} \gamma^\mu v_r c^{rz} tr_z[F_{\sigma\delta} \lambda] \\ & + \frac{1}{\sqrt{2}}x_r^m c^{rz} tr_z(\bar{\chi}^m \gamma^{\mu\nu} \lambda F_{\mu\nu}) \\ & + i v_r c^{rz} tr_z(\bar{\lambda} \gamma^\mu D_\mu \lambda) \\ & + \frac{i}{12}x_r^m x_s^m H_{\mu\nu\rho}^r c^{sz} tr_z(\bar{\lambda} \gamma^{\mu\nu\rho} \lambda) \quad , \end{aligned} \quad (18)$$

up to terms quartic in the fermions that we have omitted for simplicity. Actually, this is not quite the proper lagrangian of the theory, that would involve some peculiar couplings of the antisymmetric tensors, following the original proposal of Pasti, Sorokin and Tonin [13], necessary to enforce their (anti)self-duality conditions. However, using these only after varying, eq. (18) yields the proper equations of all fields aside from the antisymmetric tensors. Moreover, varying with respect to the antisymmetric tensors yields their second-order equations, namely the divergence of the (anti)self-duality conditions of eq. (12). The vector equation is not covariant already to lowest order, but all terms of higher order in the fermi fields do not affect the gauge anomaly.

It is now straightforward to study the effect of supersymmetry transformations, and one obtains

$$\delta B(eq. B) + \delta F(eq. F) = \mathcal{A}_\epsilon \quad , \quad (19)$$

where F and B denote collectively the fermi and bose fields, aside from the antisymmetric tensors. Moreover, in eq. (19) one has to impose the complete (anti)self-duality conditions on the 3-form field strengths. The complete supersymmetry anomaly $\mathcal{A}_\epsilon$ that solves the Wess-Zumino consistency conditions comprises eq. (16), together with the additional terms

$$\begin{aligned} \Delta \mathcal{A}_\epsilon = & \\ & ec \cdot c' tr\{\frac{i}{2}\delta_\epsilon A_\mu F_{\nu\rho}(\bar{\lambda}' \gamma^{\mu\nu\rho} \lambda') \end{aligned}$$

$$\begin{aligned}
& + \frac{i}{2} \delta_\epsilon A_\mu (\bar{\lambda} \gamma^{\mu\nu\rho} \lambda') F'_{\nu\rho} \\
& + i \delta_\epsilon A_\mu (\bar{\lambda} \gamma_\nu \lambda') F'^{\mu\nu} \\
& + \frac{1}{32} \delta_\epsilon e_\mu^a (\bar{\lambda} \gamma^{\mu\alpha\beta} \lambda) (\bar{\lambda}' \gamma_{a\alpha\beta} \lambda') \\
& - \frac{1}{2\sqrt{2}} \delta_\epsilon A_\alpha (\bar{\lambda} \gamma^\alpha \gamma^\beta \gamma^\gamma \lambda') (\bar{\lambda}' \gamma_\beta \Psi_\gamma) \\
& + \frac{x_s^m c^{sz'}}{v_t c^{tz'}} \left[-\frac{3i}{2\sqrt{2}} \delta_\epsilon A_\alpha (\bar{\lambda} \gamma^\alpha \lambda') (\bar{\lambda}' \chi^m) \right. \\
& - \frac{i}{4\sqrt{2}} \delta_\epsilon A_\alpha (\bar{\lambda} \gamma^{\alpha\beta\gamma} \lambda') (\bar{\lambda}' \gamma_\beta \chi^m) \\
& \left. - \frac{i}{2\sqrt{2}} \delta_\epsilon A_\alpha (\bar{\lambda} \gamma_\beta \lambda') (\bar{\lambda}' \gamma^{\alpha\beta} \chi^m) \right] \} \quad . \quad (20)
\end{aligned}$$

In general, the anomaly is defined up to the variation of a local functional. In our case, this reflects the freedom of adding to the lagrangian the term

$$\mathcal{L}_{\lambda^4} = e^{\frac{\alpha}{2} c_r^z c^{rz'}} tr_{z,z'} [(\bar{\lambda} \gamma^\alpha \lambda') (\bar{\lambda} \gamma_\alpha \lambda')] \quad , \quad (21)$$

that modifies the supersymmetry anomaly according to

$$\Delta' \mathcal{A}_\epsilon = \delta_\epsilon \mathcal{L}_{\lambda^4} \quad . \quad (22)$$

This term affects the field equation of the gaugino, and thus the commutator of eq. (17) has to be reconsidered in order to accommodate this modification. As a result, after using the modified field equation, one is left with the residual term

$$\begin{aligned}
& [\delta_1, \delta_2]_{extra} \lambda = \\
& \frac{c \cdot c'}{v \cdot c} tr_{z'} \left[-\frac{1}{4} (\bar{\epsilon}_1 \gamma_\alpha \lambda') (\bar{\epsilon}_2 \gamma_\beta \lambda') \gamma^{\alpha\beta} \lambda \right. \\
& - \frac{\alpha}{2} (\bar{\lambda} \gamma_\alpha \lambda') (\bar{\epsilon}_1 \gamma_\beta \lambda') \gamma^{\alpha\beta} \epsilon_2 \\
& + \frac{\alpha}{16} (\bar{\lambda} \gamma_{\alpha\beta\gamma} \lambda') (\bar{\epsilon}_1 \gamma^\gamma \lambda') \gamma^{\alpha\beta} \epsilon_2 \\
& + \frac{\alpha}{16} (\bar{\lambda} \gamma_\gamma \lambda') (\bar{\epsilon}_1 \gamma^{\alpha\beta\gamma} \lambda') \gamma_{\alpha\beta} \epsilon_2 \\
& + \frac{1-\alpha}{4} (\bar{\lambda} \gamma_\alpha \lambda') (\bar{\epsilon}_1 \gamma^\alpha \lambda') \epsilon_2 - (1 \leftrightarrow 2) \\
& \left. + \frac{1-\alpha}{16} (\bar{\epsilon}_1 \gamma^\alpha \epsilon_2) (\bar{\lambda}' \gamma_{\alpha\beta\gamma} \lambda') \gamma^{\beta\gamma} \lambda \right] \quad . \quad (23)
\end{aligned}$$

As we will see in the next section, the presence of this 2-cocycle is strictly correlated to the presence of the anomaly, and indeed one can not eliminate

it, as befits a theory that has a built-in anomaly for every value of α . Finally, one may show that the divergence of the complete gravitino equation is proportional to the complete supersymmetry anomaly.

2. Comments on Gauge and Supersymmetry Anomalies

If a supersymmetric Yang-Mills theory formulated in the Wess-Zumino gauge has a gauge anomaly, it must also have a supersymmetry anomaly, on account of the Wess-Zumino consistency conditions

$$\begin{aligned}
& \delta_\epsilon \mathcal{A}_\Lambda = \delta_\Lambda \mathcal{A}_\epsilon \quad , \\
& \delta_{\epsilon_1} \mathcal{A}_{\epsilon_2} - \delta_{\epsilon_2} \mathcal{A}_{\epsilon_1} = \mathcal{A}_{\tilde{\Lambda}} \quad , \quad (24)
\end{aligned}$$

where $\tilde{\Lambda}$ denotes the gauge parameter generated by the commutator of two supersymmetry transformations. The simplest example is provided by the globally supersymmetric Yang-Mills theory in four dimensions. From the four-dimensional gauge anomaly

$$\mathcal{A}_\Lambda^4 = tr[\Lambda(dA)^2 + \frac{i}{2} g d\Lambda A^3] \quad , \quad (25)$$

and from eqs. (24), one can determine the form of the corresponding supersymmetry anomaly [14]. In particular, from the first of eqs. (24) one obtains

$$\mathcal{A}_\epsilon^4 = tr[\delta_\epsilon A A(dA) + \delta_\epsilon A(dA)A - \frac{3ig}{2} \delta_\epsilon A A^3] \quad . \quad (26)$$

This satisfies the second of eqs. (24) if augmented by the gauge-invariant term

$$\Delta \mathcal{A}_\epsilon^4 = -\frac{i}{2} tr[\delta_\epsilon A \bar{\lambda} \gamma^{(3)} \lambda + \bar{\lambda} \delta_\epsilon A \gamma^{(3)} \lambda] \quad , \quad (27)$$

where $\gamma^{(3)} = \gamma_{\mu\nu\rho} dx^\mu \wedge dx^\nu \wedge dx^\rho$, λ is a right-handed Weyl spinor and

$$\begin{aligned}
& \delta A_\mu = \frac{i}{\sqrt{2}} (\bar{\epsilon} \gamma_\mu \lambda - \bar{\lambda} \gamma_\mu \epsilon) \quad , \\
& \delta \lambda = \frac{1}{2\sqrt{2}} F_{\mu\nu} \gamma^{\mu\nu} \epsilon \quad . \quad (28)
\end{aligned}$$

Although the algebra closes only on the field equation of λ , in four dimensions it is not possible

to generate from eqs. (24) a term proportional to $\gamma^\mu D_\mu \lambda$, and thus the Wess-Zumino consistency conditions close also off-shell. This was originally observed in [14].

On the other hand, in six dimensions, starting from the residual gauge anomaly

$$\mathcal{A}_\Lambda^6 = -c^{rz} c_r^{z'} tr_z(\Lambda dA) tr_{z'}(F^2) \quad , \quad (29)$$

the first of eqs. (24) implies a corresponding supersymmetry anomaly

$$\begin{aligned} \mathcal{A}_\epsilon^6 = & -c^{rz} c_r^{z'} [tr_z(\delta_\epsilon AA) tr_{z'}(F^2) \\ & - 2tr_z(\delta_\epsilon AF) \omega_3^z] \quad , \end{aligned} \quad (30)$$

that has to be augmented by the gauge-invariant terms

$$\begin{aligned} \Delta \mathcal{A}_\epsilon = & A c_r^z c^{rz'} tr_z[\delta_\epsilon A_\mu F_{\nu\rho}] tr_{z'}[\bar{\lambda}' \gamma^{\mu\nu\rho} \lambda'] \\ & + B c_r^z c^{rz'} tr_z[\delta_\epsilon A_\mu \bar{\lambda}] \gamma^{\mu\nu\rho} tr_{z'}[\lambda' F'_{\nu\rho}] \\ & + C c_r^z c^{rz'} tr_z[\delta_\epsilon A_\mu \bar{\lambda}] \gamma_\nu tr_{z'}[\lambda' F'^{\mu\nu}] \quad , \end{aligned} \quad (31)$$

with coefficients satisfying the relations

$$A + B = i \quad , \quad C = 4A - 2B \quad . \quad (32)$$

As pointed out in the previous Section, these leave one undetermined parameter, in agreement with the fact that the anomaly is defined up to the variation of a local functional. Indeed, the variation

$$\delta[(\bar{\lambda} \gamma^\alpha \lambda')(\bar{\lambda} \gamma_\alpha \lambda')] \quad (33)$$

yields the sum in eq. (31) with $A + B = 0$ and $C = 4A - 2B$, so that adding it to $\mathcal{A}_\epsilon$ effectively does not affect eqs. (32). The main result, anticipated in [14], is that in six dimensions the last of eqs. (24) generates terms containing one derivative and four gaugini, that cancel only using the Dirac equation $\gamma^\mu D_\mu \lambda = 0$. To reiterate, in six dimensions the Wess-Zumino consistency conditions for an anomalous Yang-Mills theory close only on shell.

The freedom of adding to the anomaly a $\delta \lambda^4$ term is another, related feature of the six-dimensional case that we would like to stress. Indeed, one can easily observe that in D dimensions the effective weight³ of the supersymmetry

³As usual in supersymmetric theories, this is 1 for derivatives, 1/2 for fermi fields, and 0 for bose fields.

anomaly is $(D-1)/2$, while that of the supersymmetry variation in (33) is 5/2. Therefore, the undetermined λ^4 contribution is present only in six dimensions. Moreover, in all cases the breaking of supersymmetry affects the algebra on the gaugino, that as usual closes only on the field equation.

All this has direct implications for six dimensional (1,0) supergravity. In this case, the Wess-Zumino conditions include additional contributions, and become

$$\begin{aligned} \delta_\epsilon \mathcal{A}_\Lambda &= \delta_\Lambda \mathcal{A}_\epsilon \quad , \\ \delta_{\epsilon_1} \mathcal{A}_{\epsilon_2} - \delta_{\epsilon_2} \mathcal{A}_{\epsilon_1} &= \mathcal{A}_{\bar{\epsilon}} + \mathcal{A}_{\bar{\Lambda}} \quad . \end{aligned} \quad (34)$$

Again, they are only satisfied on-shell and, more precisely, if $\alpha = 0$ one obtains

$$\begin{aligned} (\delta_{\epsilon_1} \mathcal{A}_{\epsilon_2} - \delta_{\epsilon_2} \mathcal{A}_{\epsilon_1})_{extra} = & c \cdot c' tr \left\{ \frac{1}{8} (\bar{\epsilon}_1 \gamma_\sigma \epsilon_2) (\bar{\lambda} \gamma_\mu \lambda') (\bar{\lambda} \gamma^\mu \gamma^\sigma [eq \cdot \lambda']_{\alpha=0}) \right. \\ & - \frac{1}{16} [\bar{\epsilon}_1 \gamma_{\sigma\delta\tau} \epsilon_2]_i \{ [\bar{\lambda} \gamma^\tau \lambda']_i (\bar{\lambda} \gamma^{\sigma\delta} [eq \cdot \lambda']_{\alpha=0}) \\ & \left. + [\bar{\lambda} \gamma^\tau \lambda]_i (\bar{\lambda}' \gamma^{\sigma\delta} [eq \cdot \lambda']_{\alpha=0}) \} \right\} \quad , \end{aligned} \quad (35)$$

where the terms within square brackets are fermionic bilinears with a non-trivial symplectic structure defined by the relation

$$[\bar{\chi} \Psi]_i = \sigma_{ia}{}^b \bar{\chi}_b \Psi^a \quad . \quad (36)$$

Still, the identity

$$\begin{aligned} & c \cdot c' tr \left\{ \frac{e}{8} (\bar{\epsilon}_1 \gamma_\sigma \epsilon_2) (\bar{\lambda} \gamma_\mu \lambda') (\bar{\lambda} \gamma^\mu \gamma^\sigma [eq \cdot \lambda']_\alpha) \right. \\ & - \frac{e}{16} [\bar{\epsilon}_1 \gamma_{\sigma\delta\tau} \epsilon_2]_i \{ [\bar{\lambda} \gamma^\tau \lambda']_i (\bar{\lambda} \gamma^{\sigma\delta} [eq \cdot \lambda']_\alpha) \\ & \left. + [\bar{\lambda} \gamma^\tau \lambda]_i (\bar{\lambda}' \gamma^{\sigma\delta} [eq \cdot \lambda']_\alpha) \} \right\} = \\ & c \cdot c' tr \left\{ \frac{e}{8} (\bar{\epsilon}_1 \gamma_\sigma \epsilon_2) (\bar{\lambda} \gamma_\mu \lambda') (\bar{\lambda} \gamma^\mu \gamma^\sigma [eq \cdot \lambda']_{\alpha=0}) \right. \\ & - \frac{e}{16} [\bar{\epsilon}_1 \gamma_{\sigma\delta\tau} \epsilon_2]_i \{ [\bar{\lambda} \gamma^\tau \lambda']_i (\bar{\lambda} \gamma^{\sigma\delta} [eq \cdot \lambda']_{\alpha=0}) \\ & \left. + [\bar{\lambda} \gamma^\tau \lambda]_i (\bar{\lambda}' \gamma^{\sigma\delta} [eq \cdot \lambda']_{\alpha=0}) \} \right\} \\ & + \frac{e\alpha}{8} \frac{c_r^z c^{rz'} c_s^{z'} c^{sz''}}{v_t c^{tzz'}} tr_{z,z',z''} [\bar{\epsilon}_1 \gamma^{\mu\nu\rho} \epsilon_2]_i [\bar{\lambda} \gamma_\mu \lambda]_i \cdot \\ & [\bar{\lambda}' \gamma_\nu \lambda']_j [\bar{\lambda}'' \gamma_\rho \lambda'']_j \end{aligned} \quad (37)$$

implies that the last term should somehow be generated in the anomaly, if the Wess-Zumino condition is to close for any value of α . In the presence of $\mathcal{L}_{\lambda^4}$, however, the anomaly is modified by eq.

(22). Using the field equation of the gaugini for arbitrary α , the last of eqs. (24) on this term yields,

$$[\delta_{\epsilon_1}, \delta_{\epsilon_2}] \mathcal{L}_{\lambda^4} = \delta_{\bar{\epsilon}} \mathcal{L}_{\lambda^4} + \delta_{\alpha} \mathcal{L}_{\lambda^4} \quad , \quad (38)$$

where δ_{α} is the transformation defined in eq. (23). The end result is that this transformation exactly equals the last term in eq. (37), and thus one can understand the rationale behind the occurrence of the 2-cocycle in the algebra on the gaugini: it lets the Wess-Zumino conditions close precisely on the field equations determined by the algebra. Since the Wess-Zumino conditions need only the equation of the gaugini, only these fields sense the additional transformation.

To summarize, in six dimensions the Green-Schwarz mechanism exhibits some relevant novelties, since the Green-Schwarz term is in this case $B \wedge F \wedge F$. Although generated by a one loop contribution from the string viewpoint, in the low-energy supergravity this introduces tree-level anomalies that are compensated by fermion loops. This fact has important consequences for the low-energy couplings, that play a central role in perturbative six-dimensional type-I vacua and in recent discussions of string duality [15], some of which we have tried to elucidate.

REFERENCES

1. M.B. Green and J.H. Schwarz, Phys. Lett. 149B (1984) 117.
2. S. Randjbar-Daemi, A. Salam, E. Sezgin and J. Strathdee, Phys. Lett. 151B (1985) 351.
3. M. Bianchi and A. Sagnotti, Phys. Lett. B247 (1990) 517; Nucl. Phys. B361 (1991) 519; E. Gimon and J. Polchinski, hep-th/9601038; A. Dabholkar and J. Park, hep-th/9602030, 9604178; E. Gimon and C.V. Johnson, hep-th/9604129; M. Berkooz, R. Leigh, J. Polchinski, J.H. Schwarz, N. Seiberg and E. Witten, Nucl. Phys. B475 (1996) 115; J. Blum and A. Zaffaroni, Phys. Lett. B387 (1996) 71; J. Blum, Nucl. Phys. B486 (1997) 34; A. Sen, Nucl. Phys. B475 (1996) 562, Phys. Rev. D55 (1997) 7345, hep-th/9709159; J. Blum and K. Intriligator, hep-th/9705030, 9705044; J. Blum, K.R. Dienes, hep-th/9707148, 9707160; K. Dasgupta and S. Mukhi, Phys. Lett. B385 (1996) 125; C. Angelantonj, M. Bianchi, G. Pradisi, A. Sagnotti and Ya.S. Stanev, Phys. Lett. B387 (1996) 743. For a review, see: L.E. Ibáñez and A.M. Uranga, hep-th/9707075.
4. A. Sagnotti, in “Non-Perturbative Quantum Field Theory”, eds. G. Mack et al (Pergamon Press, 1988), p. 521.
5. A. Sagnotti, Phys. Lett. 294B (1992) 196.
6. S. Ferrara, F. Riccioni and A. Sagnotti, hep-th/9711059.
7. L.J. Romans, Nucl. Phys. B276 (1986) 71.
8. H. Nishino and E. Sezgin, Nucl. Phys. B144 (1986) 353.
9. H. Nishino and E. Sezgin, hep-th/9703075.
10. S. Ferrara, R. Minasian and A. Sagnotti, Nucl. Phys. B474 (1996) 323.
11. J. Wess and B. Zumino, Phys. Lett. B37 (1971) 95.
12. J.H. Schwarz, Nucl. Phys. B226 (1983) 269.
13. P. Pasti, D. Sorokin, M. Tonin, Phys. Rev. D55 (1995) 6292.
14. G. Girardi, R. Grimm and R. Stora, Phys. Lett. B156 (1985) 203; L. Bonora, P. Pasti and M. Tonin, Phys. Lett. B156 (1985) 341; E. Guadagnini, K. Konishi and M. Mintchev, Phys. Lett. B157 (1985) 37; N.K. Nielsen, Nucl. Phys. B244 (1984) 499; H. Itoyama, V.P. Nair and H. Ren, Nucl. Phys. B262 (1985) 317; E. Guadagnini and M. Mintchev, Nucl. Phys. B269 (1986) 543; S. Ferrara, A. Masiero, M. Porrati and R. Stora, Nucl. Phys. B417 (1994) 238.
15. E. Witten, Nucl. Phys. B460 (1996) 541; M.J. Duff, R. Minasian and E. Witten, Nucl. Phys. B465 (1996) 413; N. Seiberg and E. Witten, Nucl. Phys. B471 (1996) 121; M.J. Duff, H. Lü and C. Pope, Phys. Lett. B378 (1996) 101.